



Hawaan Forest Conservation Trust

Monthly site inspection report:

22.04.2024

Summary:

- With the onset of autumn on the KwaZulu-Natal coastal belt, the forest has now entered its semi-deciduous phase. Unless we receive significant rainfall, we expect the forest to dry out until around mid-September or even mid-October.
- The seasonal migrant and endangered Spotted Ground Thrush has been frequently observed at the Hawaan and neighboring Sibaya Forest during April. The Song-meter used to measure bioacoustics has been in the testing phase, so we should be able to formally determine the occurrence of this bird using this tool.
- Work conducted during April by the HCT operations team has included standard operations such as mowing, pruning, and weeding of the trail network around the grassland and forest. Spot spraying of invasive and woody plant species, specifically Searsia nebulosa, has continued. We will prepare tracer belts for the prescribed burn during the month of May.
- An additional four camera traps have been set up along the southern portion of the Bulbul trail. Some of the observations included in the pages below feature two species of forest doves, the Lemon Dove and Tambourine Dove.
- Work has been conducted on the eastern (neighboring) THD/WESSA component of the forest, which has included:
 - (i) Development of trail links with the Hawaan trail network; (ii) Maintenance and redevelopment of the overgrown THD/WESSA trail network; snare patrol along the eastern M4 Boundary; removal of invasive plant species.
- The construction of a new gate at the forest management zone, and a site camp for the security contractor will require some cleaning up once this work has been completed. This will include formalizing stockpiling areas for landscaping mulch, etc.




Figure 1. Numbers on the map refer to items below.

(1)



A new gate at the forest management zone has been installed. The upgrade includes a permanent security camera and replacement of the fencing in the area.

(2)




The sign on Herald Drive needs to be straightened up. HOA have offered to help with this during the up coming fence installation.

(3)



Housekeeping issues at the Forest Management Zone: The old infrastructure of the security cameras will be removed within the next week.

(4)



Trails linking the HCT to the WEssa property have been opened up during April, including maintenance of the WEssa trail.

(5)



Flowering on the forest edge in April are *Hibiscus surattensis*, which is quite a prickly and scandent climbing shrub.

(6)



The central link Bushshrike trail link to WEssa forest area has also been opened up the pipeline trail.

(7)



(8)



The Solitary Bees have just about gone into winter diapause so the forest edge trail on the corner of the Bushshrike will be cut by the end of April.

Flowering on the forest edge during April are African Blue Bell, *Wahlenbergia undulata*.

(9)



(10)



Monthly management of *Eugenia capensis* and *Allophylus natalensis* has been conducted in grassland Block 3.

Grassland trails were cut in late April and were looking neat during the inspection.

(11)



(12)



Wild Lions Paw have done well this summer and have continued to flower over late March and into April.

Solanum seaforthianum on the Bush Shrike forest trail are to be removed in April.

(13)



Camera traps and water points have been serviced in April. The new camera's have replaced the older model camera which were previously located at this water point.

(15)



Bernard, and the WESSA team have now cleared links and have opened up the WESSA loop trail during April.

(17)



Bernard, and the WESSA team have now cleared links and have opened the WESSA loop trail during April. Also feeding off *Allophylus natalensis* during April are Pearl Charaxes butterflies

(14)



The conservation team to rake pathway to the Bush Shrike trail before the end of April.

(16)




One of few plants flowering in the forest during April are *Allophylus natalensis*, these plants are alive with insect activity, including this striped eyed Lagoon Fly.

(18)



Yellow-collared carpenter, *Xylocarpa flavidollis*, was also noted feeding off the nectar on the *Tephrosia*'s within the grassland.

Camera Trap imagery from April 2024



Vervet monkeys have been enjoying the fresh water provided HCT at the Bushshrike water point



As have many of the Red Duiker



The large male bush-buck



and the beautiful female bush-buck, all look in fine condition.



Two beautiful forest doves, the rarer forest, Lemon Dove..



and the more common, more often associated with woodlands, Tambourine Dove, were captured on the camera during April

Operations: Tasks 2024

	3.7	Keep Tilley Huts neat and tidy sweep daily, clean windows every 10 days check maintenance and rusting on a monthly basis and report to HCT members.	Weekly	x	x	z	x	-	-	-	-	-	-
	3.8	Check that trail cameras y ggm"to ensure they are not stolen.	Daily	x	x	z	x	-	-	-	-	-	-
	3.9	Cleaning of shower on a daily basis after use.	na	-	-	z	x	-	-	-	-	-	-
	3.10	Check and order cleaning consumables for shower and toilet	na	-	-	-	-	-	-	-	-	-	-
Woody plant management	4.1	On-going daily work is clearing invasive introduced plants such as Pepper Trees, Triffid etc..in both the forest and grassland systems.	Weekly	x	x	z	x	-	-	-	-	-	-
		Checking for regrowth of IAPs monthly and set out new work areas each month.	Monthly	x	x	-	x	-	-	-	-	-	-
	4.2	Removal of selected indigenous woody encroachment species such Silver Oak in the grassland section	Monthly	-	x	-	x	-	-	-	-	-	-
	4.3	Monitoring herbicide stores on a monthly	Monthly	x	x	-	x	-	-	-	-	-	-
Fire management	5.1	Ensure tracers belts and pathways in the grassland open before prescribed burn	Yearly	-	-	-	-	-	-	-	-	-	-
	5.2	Service fire-fighting equipment once before the annual burn is completed and then oil up so that ne-t season the equipment has not seized up with rust. <i>Fire fighting</i> =equipment one drip torch, 3 rubber beaters with handles, two Knapsack sprayers.	Yearly	-	-	-	-	-	-	-	-	-	-
Consumables	6.1	Order uniforms and personal protective gear for the 2 staff.	Yearly	-	-	-	-	-	-	-	-	-	-

Item	Herbicide name	Active ingredient	Type	Litres in stock	Date used	Species applied	Location	Active ingredient	Type	Litres in stock	Date used	Species applied	Location	
1.	Gladiator, pre-mix	Picloram	Selective	15L (pre-mix)	16/4/22	Creeping inch	Forest mgmt zone	5.	Plenum	Picloram	Selective	18L	13/4	Pepper, Yellowbe
2.	Round-up	Glyphosate	Non-selective	3L	-	-	-	6.	Plenum	Picloram	Selective	5L	05/9	IlS Pepper,
3.	Gladiator, premix	Picloram	Selective	Used 15L	15/11/22	Brazilian pepper	Eastern grassland	7.	Plenum	Picloram	Selective	5L	05/03/24	Searia
4.	Plenum 160	Picloram	Selective	20L – concentrate.	25/02/23	Brazilian pepper – Yellow bellis.	Eastern grassland	8.						

Observations from the camera traps in February showed the usual allotment of animals using the water provided by the HCT staff to bathe and drink. We are working towards formalising the logging of this data on a consistent basis so that it can be used as baseline information for the forest and to track change.



Spotted Genet



Tambourine Dove




Red Duiker



Blue Duiker



Spotted Eagle Owl



Water Mongoose

Camera Trap highlights from November 2023. Hawaan Forest.



Large Spotted Genet



Female Bushbuck



Red Capped Robin Chat



Pair of Red Duikers



Purple crested turaco




Tambourine dove

The forest fauna have been out in full force during August and the action has clearly revolved around the water point at the entrance to the Bush Shrike trail. One again some beautiful forest images have been captured.



A lemon dove and male blue duiker



A Red capped robin chat with a Olive thrush



Juvenile crowned eagle



A family of bushbuck




Large spotted Genet



Dwarf mongoose

Some beautiful camera trap images were taken at the forest water point during July, 2022.



Red Duiker, female.



Bushbuck females.



Crested guinea fowl.



Red Duiker, male



Vervet Monkey, Banded Mongoose, and Guinea fowl.



Banded Mongoose.

Some beautiful camera trap images were taken within the dry forest interior during June.



Bush buck female)



Bush buck (male)



Bush buck (female)



Red duiker (female)



Bush buck (female)



Blue duiker (pair)

Camera Trap imagery from March 2024



The have been some great mixed foraging parties captured on the camera's during March. Here, Crested Guinea fowls and Red Duiker.



On the Bush-Shrike trail, Crested Guinea fowl and Bush Buck.




The recent hot weather has prompted a diverse array animals captured at the water-hole on the Bush-shrike trail. Such as the Crowned eagle drinking at the water-hole.



A family of Purple Crested Turaco's.



A nice large Bushbuck passing by the water-hole on the Bush Shrike trail.




A small, blue Duiker close up at the water-hole.

The forest fauna have been out in full force during August and the action has clearly revolved around the water point at the entrance to the Bush Shrike trail. One again some beautiful forest images have been captured.



A lemon dove and male blue duiker



A Red capped robin chat with a Olive thrush



Juvenile crowned eagle



A family of bushbuck




Large spotted Genet



Dwarf mongoose

Some beautiful camera trap images were taken at the forest water point during July, 2022.



Red Duiker, female.



Bushbuck females.



Crested guinea fowl.



Red Duiker, male



Vervet Monkey, Banded Mongoose, and Guinea fowl.



Banded Mongoose.

Some beautiful camera trap images were taken at the forest water point during July, 2022.



Red Duiker, female.



Bushbuck females.




Crested guinea fowl.



Red Duiker, male




Vervet Monkey, Banded Mongoose, and Guinea fowl.



Banded Mongoose.

September 2022

The Red Duiker are the most beautiful forest antelope we have at the Hawaan Forest. Larger, and less common than their smaller cousins the Blue Duiker, they may reach up to 12 kg. They are coloured a deep chestnut red which contrasts pleasingly with the luminescent greens of the forest foliage in summer and similarly with the dry-scrub brown of the deciduous winter phase of the forest. These camera trap images, taken this month in September, showcase this most graceful forest antelope amongst the soft greens and browns of the forest before the onset of spring.



Camera trap images February 2022

During Jan/Feb the HCT have worked closely with the HRA and Marshall security. The camera trap maintenance and daily checking is being conducted by Marshall security, though the HCT still have camera located in locations that will capture beautiful and interesting wildlife images. One highlight from February was a Bushbuck male captured at dawn on the interface between the estate and the grassland.



Bushbuck male at dawn.




Crested-guineafowl in the forest at night!



Blue duiker in the forest



Lemon dove in the forest leaf litter



The pair of resident water-mongoose



A pair of Olive-thrush's


Camera trap observation Jan 2022. Due to the heat in January the most notable camera trap observations for Jan 2021 focused on the water trough at Bushshrike trail. An exciting observation was a juvenile crowned eagle having a drink and cooling off in the water.



Crested Guinea fowl



Purple crested turaco



Crowned eagle (Juvenile)



Female bushbuck



Wood owl



Tambourine dove

Six of the seven camera have been in operation during November-December. The Seventh camera is due to be installed within the next week. Camera batteries appear to last between 3-4 four weeks, while each camera used between 500 mb and 1gig of data. The data budget for each camera should be in the order of R 50-75 per month. Importantly no non-resident 'humans' have been noted on the cameras. AS will continue to monitor the camera in December but the way forward in January will be garner support from the HOA security to help with monitoring. Some interesting new species were observed over the Nov/Dec period, for example: on Camera V, a Steppe Eagle. Other observations in the images are : i) = Large spotted Genet ; ii) & vii) Female bushbuck; i) Water mongoose i) Male bushbuck.



⌚ v)



⌚ i)



⌚ iii)



⌚ vii)





⌚ i)



⌚ j)

Camera trap images October-November 2021. New species observations in November were Water Mongoose. Other sightings include: Bushbuck family activity, Blue Duiker pairs and Crested Guine fowl.



Camera trap images March 2022

During March the HCT have again worked closely with the HRA and Marshall security. Highlights from March 2022 include: Many different forest fauna utilising the water point at the BushShrike junction, including: Blue Duiker; Black Sparrow Hawk; Purple Turaco; and a Wood Owl. The camera on the Bush-Shrike trail also picked up some beautiful images of foraging Banded Mongoose and Crested Guinea fowl.



Blue Duiker



Black Sparrow-Hawk



Purple Turaco



Wood Owl




Banded Mongoose



Crested Guinea fowl

Camera trap images September /October 2021

Given that we are still working out how best to deploy the camera traps in the nature reserve, some of the images received so far have been beautiful. Species noted below are as follows: Crested guinea fowl, spotted Genet, and female, male and juvenile bushbuck. Other forest antelope noted have been both red and blue duiker. Four cameras have now been deployed. Their locations have been sited in Fig. 1.



Some beautiful camera trap images were taken within the dry forest interior during June.



Bush buck female)



Bush buck (male)



Bush buck (female)



Red duiker (female)



Bush buck (female)



Blue duiker (pair)

Sewer break: During the rainfall and flooding in mid-April the municipal sewer pipeline on the boundary of the HFT property ruptured. The damage is therefore not in the HFT property but will effect the species in forest. The event resulted in a high pressure flow of sewerage which eroded sections of the pipeline infrastructure while causing a plume of waste into the forest. Fortunately, it appears that the runoff did not undercut the pipeline too badly and good work from the city ensured that the issue was resolved timorously. What remains, is for the city to repair the ailing infrastructure and to co-ordinate a clean-up of the litter/plastics which was discharged into the forest.



Fig 1a. The flow of the effluent took a north-easterly direction.



Fig. 2b. Some areas of the pipeline have been undercut and will need to be repaired.



Fig 2c. The manholes on the pipeline remain open for the time being.




Fig2d. Some of the surrounding vegetation at the source point has collapsed but damage does not seem to be extensive. However, a full inspection has not been conducted.



Fig. 2e. The sewage plume continues in a north-east direction down the forest slope. Litter from this plume should be cleaned-up.

Some beautiful images and interesting sighting on the camera traps were observed during May. Firstly, a lovely bushbuck male on the Bushshrike trail camera. While many birds (tambourine dove, dark backed weaver, natal-robin, kurrichane thrush, wood owl, purple-headed turaco, thick-billed weaver and ashy fly catchers) and forest mammals (red and blue duikers, bushbuck, water mongoose, vervet monkey) have also utilised the water point.




Male bushbuck




Tambourine dove




Blue duiker (female)



Dark backed forest weaver



Purple-headed turaco



Vervet monkey

The most interesting camera trap images for October 2022 were of Bush buck males rutting, Red duiker and parties of Crested guinea fowl. Unfortunately we are yet to pick up Bushpig in the forest.



Bushbuck rutting



Bushbuck rutting



Crested Guinea fowl



Crested Guinea fowl



Red duiker



Red duiker

September 2022

The Red Duiker are the most beautiful forest antelope we have at the Hawaan Forest. Larger, and less common than their smaller cousins the Blue Duiker, they may reach up to 12 kg. They are coloured a deep chestnut red which contrasts pleasingly with the luminescent greens of the forest foliage in summer and similarly with the dry-scrub brown of the deciduous winter phase of the forest. These camera trap images, taken this month in September, showcase this most graceful forest antelope amongst the soft greens and browns of the forest before the onset of spring.



Camera trap images February 2022

During Jan/Feb the HCT have worked closely with the HRA and Marshall security. The camera trap maintenance and daily checking is being conducted by Marshall security, though the HCT still have camera located in locations that will capture beautiful and interesting wildlife images. One highlight from February was a Bushbuck male captured at dawn on the interface between the estate and the grassland.



Bushbuck male at dawn.




Crested-guineafowl in the forest at night!



Blue duiker in the forest



Lemon dove in the forest leaf litter



The pair of resident water-mongoose



A pair of Olive-thrush's


Camera trap observation Jan 2022. Due to the heat in January the most notable camera trap observations for Jan 2021 focused on the water trough at Bushshrike trail. An exciting observation was a juvenile crowned eagle having a drink and cooling off in the water.



Crested Guinea fowl



Purple crested turaco



Crowned eagle (Juvenile)



Female bushbuck



Wood owl



Tambourine dove

Six of the seven camera have been in operation during November-December. The Seventh camera is due to be installed within the next week. Camera batteries appear to last between 3-4 four weeks, while each camera used between 500 mb and 1gig of data. The data budget for each camera should be in the order of R 50-75 per month. Importantly no non-resident 'humans' have been noted on the cameras. AS will continue to monitor the camera in December but the way forward in January will be garner support from the HOA security to help with monitoring. Some interesting new species were observed over the Nov/Dec period, for example: on Camera V, a Steppe Eagle. Other observations in the images are : i) = Large spotted Genet ; ii) & vii) Female bushbuck; i) Water mongoose i) Male bushbuck.



⌚ v)



⌚ i)



⌚ iii)



⌚ vii)





⌚ i)



⌚ j)

Camera trap images October-November 2021. New species observations in November were Water Mongoose. Other sightings include: Bushbuck family activity, Blue Duiker pairs and Crested Guine fowl.



Camera trap images March 2022

During March the HCT have again worked closely with the HRA and Marshall security. Highlights from March 2022 include: Many different forest fauna utilising the water point at the BushShrike junction, including: Blue Duiker; Black Sparrow Hawk; Purple Turaco; and a Wood Owl. The camera on the Bush-Shrike trail also picked up some beautiful images of foraging Banded Mongoose and Crested Guinea fowl.



Blue Duiker



Black Sparrow-Hawk



Purple Turaco



Wood Owl





Banded Mongoose



Crested Guinea fowl

Camera trap images September /October 2021

Given that we are still working out how best to deploy the camera traps in the nature reserve, some of the images received so far have been beautiful. Species noted below are as follows: Crested guinea fowl, spotted Genet, and female, male and juvenile bushbuck. Other forest antelope noted have been both red and blue duiker. Four cameras have now been deployed. Their locations have been sited in Fig. 1.



Embankment slip: The extensive rain in mid-April and flooding of the Umhlanga river has triggered a slip of a steep section of the northern forest embankment (Fig. 1a,b). This section of forest appears to have been compromised for some time as exposed sections vegetation (sandy areas in Fig 1c) are visible in the orthophoto which was taken in 2018-2019. The vegetation cover on this slope prior to the slip was largely composed of pioneer vegetation such as *Chromolaena odorata* and *Brachyleana discolor*. It is not immediately clear what management or rehabilitation actions can be conducted to secure such as steep slope. It's likely that the dune slope will find a natural gradient. Aside from engineering options, one approach could be to plant or reseed this bank with fast growing and deep rooting indigenous trees such as *Albizia adianthifolia*. These will take 10 years to become established but if so they will offer resilience to this area for next decadal-scale rainfall event.



Fig. 1a. A perspective of the slip, looking north towards the bend in the Umhlanga river.



Fig. 1b. A perspective of the slip, looking south across the Umhlanga river towards the forest embankment.



Fig. 1c. An aerial perspective of the area taken in 2018/2019 shows a previous slip which would have likely occurred due to a similar rainfall or disturbance event. The erosion site is located on the outer-apex of the river bend so this may be part of ongoing catchment-scale processes.

The Polyphagous Shot Hole Borer (PHSB) is an ambrosia beetle native to Southeast Asia. In 2017 this pest was detected on London

Plane trees in the KwaZulu-Natal National Botanical Gardens, Pietermaritzburg. Its presence has since been confirmed in multiple

locations in eight provinces in South Africa. The beetle has a symbiotic relationship with the fungus *Fusarium euwallaceae*, which serves as a food source for the adults and their larvae. In susceptible trees the fungus causes a disease called Fusarium dieback, which can

lead to dying branches and tree death. The beetles attack a wide range of exotic and indigenous trees in urban, agricultural and natural landscapes.



Above: An adult female is 1.8-2.6mm long. Males are smaller and cannot fly.

PSHB is not able to complete its life cycle on all the tree species it attacks. Trees in which the beetle is able to breed and multiply are referred to as '**reproductive host trees**'. Important reproductive hosts include species of oaks, maples, willows and coral trees, avocado and castor bean. '**Non-reproductive host trees**' are attacked by the beetle, but the beetles do not establish galleries (tunnels) or breed. The fungus may, or may not cause disease. Trees are generally not expected to die. An updated list of confirmed hosts in South Africa can be viewed at www.fabinet.up.ac.za/pshb/

The movement of infested wood is an important means of spread of the beetle. Therefore, appropriate disposal of infested trees (by chipping/composting, solarization or burning) will be key to reducing the spread of this damaging pest. Surveys to monitor the spread of the beetle in South Africa are continuing. The public can assist by looking out for symptoms. Suspected instances can be reported to pshb@fabi.up.ac.za


Left: Chinese maple tree killed by PSHB and its fungus



Reproductive galleries in pecan



PSHB galleries in coral tree



Shot gun-like symptoms on London Plane

Compiled by Z.W. de Beer & T. Paap (Version 2021-03-04) www.fabinet.up.ac.za/pshb

